

EXTENSIONS OF SIMPLE MODULES FOR THE WITT ALGEBRA

KHALID RIAN

ABSTRACT. The irreducible representations of the Witt algebra W are completely known. A classification of the irreducible $U_\chi(W)$ -modules was first established by Chang and later simplified by Strade. The aim of this article is to give a classification of the extensions of the simple $U_\chi(W)$ -modules having p -character χ of height $-1, 0, 1$ and $p-1$ where p denotes the characteristic of the ground field.

After completion of this paper, it has been brought to my attention that similar results have appeared in [1]. However, the methods appearing in [1] are different from those used here.

1. INTRODUCTION

1.1. Notation. Let K be an algebraically closed field of characteristic $p > 3$ and let W be the p -dimensional $\mathbb{Z}$ -graded restricted simple Witt algebra with basis $e_{-1}, e_0, \dots, e_{p-2}$. The Lie bracket is given by

$$[e_i, e_j] = (j - i)e_{i+j} \quad \text{for all} \quad -1 \leq i, j \leq p-2,$$

where by definition $e_i = 0$ for $i \notin \{-1, 0, \dots, p-2\}$. The p -mapping is given by

$$e_i^{[p]} = \delta_{i0}e_i \quad \text{for all} \quad -1 \leq i \leq p-2.$$

For each $-1 \leq i \leq p-1$ we set

$$W_i = Ke_i \oplus \dots \oplus Ke_{p-2}$$

and we define the height $r(\chi)$ of a functional $\chi \in W^*$ to be

$$r(\chi) = \min\{i \mid -1 \leq i \leq p-1 \text{ and } \chi|_{W_i} = 0\}.$$

1.2. Simple modules. The simple W -modules were determined by Chang in [2]. We describe here very briefly the simple W -modules with a p -character χ of height $-1, 0, 1$ and $p-1$.

The author wishes to express his gratitude to Professor Jens Carsten Jantzen for his guidance and help.

1.2.1. Suppose that $\chi \in W^*$ has height at most 1. For each weight $\lambda \in \Lambda(\chi) = \{\lambda \in K \mid \lambda^p - \lambda = \chi(e_0)^p\}$ consider the χ -reduced Verma module $V_\chi(\lambda) = U_\chi(W) \otimes_{U_\chi(W_0)} K_\lambda$ where K_λ denotes K considered as a $U_\chi(W_0)$ -module via

$$e_0 \cdot 1 = \lambda \cdot 1 \quad \text{and} \quad e_i \cdot 1 = 0 \quad \text{for all } i \neq 0.$$

Theorem. *The following hold*

- (1) *If χ has height -1 , then there are p isomorphism classes of simple $U_\chi(W)$ -modules. These modules are represented by the one-dimensional trivial W -module K , the $(p-1)$ -dimensional module $S = V_\chi(p-1)/\text{Soc}_{U_\chi(W)} V_\chi(p-1)$ and the p -dimensional modules $V_\chi(\lambda)$ for $\lambda \in \{1, 2, \dots, p-2\}$.*
- (2) *If χ has height 0 , then there are $p-1$ isomorphism classes of simple $U_\chi(W)$ -modules each of dimension p and represented by $V_\chi(\lambda)$ for $\lambda \in \{0, 1, \dots, p-2\}$.*
- (3) *If χ has height 1 , then there are p isomorphism classes of simple $U_\chi(W)$ -modules each of dimension p and represented by $V_\chi(\lambda)$ for $\lambda \in \Lambda(\chi)$.*

For each $\mu \in \mathbb{F}_p = \mathbb{Z}/p\mathbb{Z}$ we let $[\mu] \in \{0, 1, \dots, p-1\}$ denote the unique representative of μ . Note that $\lambda - \lambda' \in \mathbb{F}_p$ for all $\lambda, \lambda' \in \Lambda(\chi)$. Furthermore, if χ has height 1 then $\lambda + \lambda' \notin \mathbb{F}_p$ since otherwise this would imply $\lambda, \lambda' \in \mathbb{F}_p$.

1.2.2. If $\chi \in W^*$ has height $p-1$ then every simple $U_\chi(W)$ -module with one exception is projective. The remaining simple module L has a projective cover with two composition factors both isomorphic to L , see [3, Theorem 2.6].

Remark. If $1 < r(\chi) < p-1$ then there exists a unique simple $U_\chi(W)$ -module up to isomorphism [2, Hauptsatz 1]. This module has a non-trivial self-extension.

2. EXTENSIONS OF THE REDUCED VERMA MODULES

2.1. Let $\chi \in W^*$ with $r(\chi) = -1, 0, 1$ and let $\lambda, \lambda' \in \Lambda(\chi)$. The isomorphism

$$\text{Ext}_{U_\chi(W)}(V_\chi(\lambda'), V_\chi(\lambda)) \simeq \text{Ext}_{U_\chi(W_0)}(K_{\lambda'}, V_\chi(\lambda))$$

reduces the problem of classifying the extensions of Verma modules to that of classifying the extensions of $K_{\lambda'}$ by $V_\chi(\lambda)$.

2.2. Consider a short exact sequence of $U_\chi(W_0)$ -modules

$$(1) \quad 0 \rightarrow V_\chi(\lambda) \xrightarrow{f} M \xrightarrow{g} K_{\lambda'} \rightarrow 0.$$

The Verma module $V_\chi(\lambda)$ has a basis $v_0, v_1, \dots, v_{p-1}$ given by $v_i = e_{-1}^i \otimes 1$. Set $w_i = f(v_i)$ for all i . Since $U_\chi(Ke_0)$ is a semisimple

algebra, there exists $w' \in M$ such that $e_0 w' = \lambda' w'$ and $g(w') = 1$. Thus we obtain a basis for M

$$(2) \quad w_0, w_1, \dots, w_{p-1}, w',$$

such that for all $0 \leq j \leq p-2$

$$(3) \quad e_j w_i = \begin{cases} (-1)^j \frac{i!}{(i-j)!} ((j+1)\lambda - i + j) w_{i-j}, & \text{if } j \leq i, \\ 0, & \text{otherwise.} \end{cases}$$

Furthermore, if $1 \leq j \leq p-2$ then

$$e_0 e_j w' = (e_j e_0 + j e_j) w' = (\lambda' + j) e_j w',$$

hence $e_j w'$ belongs to the 1-dimensional weight space $M_{\lambda'+j}$ spanned by $w_{[\lambda-\lambda'-j]}$. It follows that

$$(4) \quad e_j w' = \mathbf{a}_j w_{[\lambda-\lambda'-j]} \quad \text{for some } \mathbf{a}_j \in K.$$

2.3. The w' from the previous section is not unique. Since the λ' weight space in M is spanned by w' and $w_{[\lambda-\lambda']}$ any different choice for w' has the form $w' + b w_{[\lambda-\lambda']}$ for some $b \in K$. Obviously, this leads to the same $e_j w'$ if $e_j w_{[\lambda-\lambda]} = 0$. Thus, if $\lambda = \lambda'$ then $(\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_{p-2})$ is determined by the extension (1). Furthermore, we have

Lemma (A). *Let $\lambda, \lambda' \in \Lambda(\chi)$. If $(\lambda, \lambda') = (0, p-1)$ or $(\lambda', \lambda) = (p-1, 0)$ then $e_j w_{[\lambda-\lambda']} = 0$ for all $j \geq 1$.*

Note that the assumptions of the lemma imply $r(\chi) < 1$ as $\lambda, \lambda' \notin \mathbb{F}_p$ for $r(\chi) = 1$. We let $\Theta(\chi) \subset \Lambda(\chi) \times \Lambda(\chi)$ denote the subset given by

$$(\{(0, p-1), (p-1, 0)\} \cap \Lambda(\chi) \times \Lambda(\chi)) \cup \{(\mu, \mu) \mid \mu \in \Lambda(\chi)\}.$$

Lemma A and the discussion at the beginning of this section show that all the $\mathbf{a}_j$ are determined by M if $(\lambda, \lambda') \in \Theta(\chi)$. This is not the case for $(\lambda, \lambda') \notin \Theta(\chi)$. However, we have

Lemma (B). *Let $\lambda, \lambda' \in \Lambda(\chi)$ such that $(\lambda, \lambda') \notin \Theta(\chi)$. If furthermore $\lambda + \lambda' \neq p-1$ (resp. $\lambda + \lambda' = p-1$) then there is a unique choice for w' such that $e_1 w' = 0$ (resp. $e_2 w' = 0$).*

The proof is straightforward, given the fact that $e_1 w_{[\lambda-\lambda']} \neq 0$ (resp. $e_2 w_{[\lambda-\lambda']} \neq 0$) if $\lambda + \lambda' \neq p-1$ (resp. $\lambda + \lambda' = p-1$).

Remark. For each pair $(\lambda, \lambda') \notin \Theta(\chi)$, we will always assume that w' is the unique choice from the lemma. It follows, then, that the tuple $(\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_{p-2})$ determines the extension (1). We call this tuple *the $\mathbf{a}$ -datum of (1)*.

2.4. Consider now a second extension

$$(5) \quad 0 \rightarrow V_\chi(\lambda) \xrightarrow{f'} M' \xrightarrow{g'} K_{\lambda'} \rightarrow 0$$

of $U_\chi(W_0)$ -modules. If (1) and (5) are equivalent extensions then there is an isomorphism $h : M \xrightarrow{\sim} M'$ compatible with the identities in both $V_\chi(\lambda)$ and $K_{\lambda'}$

$$\begin{array}{ccccccccc} 0 & \longrightarrow & V_\chi(\lambda) & \xrightarrow{f} & M & \xrightarrow{g} & K_{\lambda'} & \longrightarrow & 0 \\ & & \parallel & & \downarrow h & & \parallel & & \\ 0 & \longrightarrow & V_\chi(\lambda) & \xrightarrow{f'} & M' & \xrightarrow{g'} & K_{\lambda'} & \longrightarrow & 0. \end{array}$$

We can choose $w'' = h(w')$ to be the analogue to w' . Then $e_i w'' = h(e_i w') = h(f(\mathbf{a}_i v_{[\lambda - \lambda' - i]})) = f'(\mathbf{a}_i v_{[\lambda - \lambda' - i]})$ for all $1 \leq i \leq p-2$. Hence $f^{-1}(e_i w')$ depends only on the class of the extension (1) and we obtain a well-defined map

$$(6) \quad \Phi_{\lambda, \lambda'}^i : \text{Ext}_{U_\chi(W_0)}(K_{\lambda'}, V_\chi(\lambda)) \rightarrow K,$$

that sends the class of (1) to $\mathbf{a}_i$. One checks that $\Phi_{\lambda, \lambda'}^i$ is linear. In Section 2.5 we show that every $\mathbf{a}_i$ with $i \geq 3$ can be expressed as a linear combination of $\mathbf{a}_1$ and $\mathbf{a}_2$. It follows that $\Phi_{\lambda, \lambda'} = (\Phi_{\lambda, \lambda'}^1, \Phi_{\lambda, \lambda'}^2)$ maps $\text{Ext}_{U_\chi(W_0)}(K_{\lambda'}, V_\chi(\lambda))$ injectively into K^2 .

Remark. If $(\lambda, \lambda') \notin \Theta(\chi)$ then $\dim \text{Ext}_{U_\chi(W_0)}(K_{\lambda'}, V_\chi(\lambda)) \leq 1$. This is a consequence of Lemma 2.3 B.

2.5. Since M is a $U_\chi(W_0)$ -module, we have

$$[e_i, e_j]w' = (e_i e_j - e_j e_i)w' \quad \text{for all } i \text{ and } j.$$

We will need the full strength of this formula later, but for now we are content to remark that in the case where $i = 1$ and $2 \leq j \leq p-3$, it yields

$$\begin{aligned} (j-1)\mathbf{a}_{j+1} = & -(-1)^j \left(\prod_{k=1}^j (\lambda - \lambda' - k) \right) (j(\lambda + 1) + \lambda' + 1)\mathbf{a}_1 \\ & - (\lambda - \lambda' - j)(\lambda + \lambda' + j + 1)\mathbf{a}_j, \end{aligned}$$

or, equivalently, by induction

$$(7) \quad \mathbf{a}_j = A_j \mathbf{a}_1 + B_j \mathbf{a}_2 \quad \text{for all } 3 \leq j \leq p-2,$$

where

$$\begin{aligned} A_j = & \frac{(-1)^j}{j-2} \left(\prod_{k=1}^{j-1} (\lambda - \lambda' - k) \right) \left((j-1)\lambda + \lambda' + j + \sum_{k=4}^j \frac{(j-k)!}{(j-3)!} \right. \\ & \left. \cdot ((j+2-k)\lambda + \lambda' + (j+3-k)) \prod_{l=0}^{k-4} (\lambda + \lambda' + j - l) \right), \end{aligned}$$

(the summation $\sum_{k=4}^j$ is understood to be 0 when $j = 3$) and

$$B_j = \frac{(-1)^j}{(j-2)!} \prod_{k=2}^{j-1} (\lambda - \lambda' - k)(\lambda + \lambda' + k + 1).$$

Consequently, all the $\mathbf{a}_i$ with $i \geq 3$ are determined by $\mathbf{a}_1$ and $\mathbf{a}_2$.

2.6. Let $\lambda, \lambda' \in \Lambda(\chi)$. We want to describe all possible $\mathbf{a}$ -data of extensions as in (1). To this end, consider an arbitrary pair $(\mathbf{a}_1, \mathbf{a}_2) \in K^2$ and use (7) to extend it to a tuple $\mathbf{a} = (\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_{p-2})$ in K^{p-2} . In order to simplify notation set $\mathbf{a}_i = 0$ for all $i > p-2$. Consider a vector space $M_{\mathbf{a}}$ with a basis $w_0, w_1, \dots, w_{p-1}, w'$ and define endomorphisms $E_j \in \text{End}_K(M_{\mathbf{a}})$ for $0 \leq j \leq p-2$ such that

$$E_j w_i = \begin{cases} (-1)^j \frac{i!}{(i-j)!} ((j+1)\lambda - i + j) w_{i-j}, & \text{if } j \leq i, \\ 0, & \text{otherwise,} \end{cases}$$

and

$$E_j w' = \begin{cases} \lambda' w', & \text{if } j = 0, \\ \mathbf{a}_j w_{[\lambda - \lambda' - j]}, & \text{otherwise.} \end{cases}$$

Furthermore we set $E_j = 0$ for $j \notin \{0, 1, \dots, p-2\}$. A comparison with the formulas in Section 2.2 shows that $\mathbf{a}$ is the $\mathbf{a}$ -datum of an extension as in (1) if and only if $M_{\mathbf{a}}$ has a W_0 -module structure such that $e_j w = E_j w$ for all $w \in M_{\mathbf{a}}$ and all $j \geq 0$ and if this W_0 -module then has p -character χ . We have on $M_{\mathbf{a}}$ a structure as W_0 -module if and only if

$$(8) \quad [E_i, E_j]w = (j-i)E_{i+j}w \quad \text{for all } i, j \geq 0 \text{ and } w \in M_{\mathbf{a}}.$$

It suffices to check this for all w in our basis. The linear map $f: V_{\chi}(\lambda) \rightarrow M_{\mathbf{a}}$ with $f(v_i) = w_i$ for all i satisfies $E_j f(v) = f(E_j v)$ for all $v \in V_{\chi}(\lambda)$ and all j . Consequently, (8) holds for all w_i and $M_{\mathbf{a}}$ has the desired structure if and only if

$$(9) \quad [E_i, E_j]w' = (j-i)E_{i+j}w' \quad \text{for all } i, j \geq 0.$$

2.7. One verifies easily that (9) holds for $i = 0$. It also holds for $i = j$. For $i = 1$ and $2 \leq j \leq p-3$ it is equivalent to (7) which holds by the definition of the $\mathbf{a}_j$ ($j \geq 3$). Furthermore, for every (i, j) in

$$\{(1, p-2), (p-2, 1)\} \cup \{(a, b) \in \mathbb{Z}^2 \mid a \neq b \text{ and } 2 \leq a, b \leq p-2\},$$

formula (9) is equivalent to the following conditions

If $i > [\lambda - \lambda' - j]$ and $j > [\lambda - \lambda' - i]$, then

$$(10) \quad \mathbf{a}_{i+j} = 0.$$

If $i > [\lambda - \lambda' - j]$ and $j \leq [\lambda - \lambda' - i]$, then

$$(11) \quad (j-i)\mathbf{a}_{i+j} = (-1)^{j+1} \frac{[\lambda - \lambda' - i]!}{([\lambda - \lambda' - i] - j)!} (j(\lambda + 1) + \lambda' + i)\mathbf{a}_i.$$

If $i \leq [\lambda - \lambda' - j]$ and $j \leq [\lambda - \lambda' - i]$, then

$$(12) \quad (j-i)\mathbf{a}_{i+j} = (-1)^i \frac{[\lambda - \lambda' - j]!}{([\lambda - \lambda' - j] - i)!} (i(\lambda + 1) + \lambda' + j)\mathbf{a}_j \\ - (-1)^j \frac{[\lambda - \lambda' - i]!}{([\lambda - \lambda' - i] - j)!} (j(\lambda + 1) + \lambda' + i)\mathbf{a}_i.$$

2.8. Suppose now that $\mathbf{a}$ satisfies (10)–(12) and hence that $M_{\mathbf{a}}$ is a W_0 -module with each e_i acting as E_i . We have then a short exact sequence

$$(13) \quad 0 \rightarrow V_{\chi}(\lambda) \xrightarrow{f} M_{\mathbf{a}} \xrightarrow{g} K_{\lambda'} \rightarrow 0.$$

Evidently, $M_{\mathbf{a}}$ has p -character χ if and only if $e_i^p w' = 0$ for all $i \geq 1$, or equivalently, if $\mathbf{a}_i e_i^{p-1} w_{[\lambda - \lambda' - i]} = 0$ for all $i \geq 1$. We have

$$\mathbf{a}_1 e_1^{p-1} w_{p-1} = -\mathbf{a}_1 \left(\prod_{j=0}^{p-2} (2\lambda - j) \right) w_0.$$

Observe that the term inside the bracket does not vanish if the height of χ is 1 because $\lambda \notin \mathbb{F}_p$ in this case. Since furthermore $e_i^{p-1} w_{[\lambda - \lambda' - i]} = 0$ if $i > 1$ or $\lambda \neq \lambda'$ we obtain the following lemma

Lemma. *$M_{\mathbf{a}}$ has p -character χ if and only if one of the following conditions holds*

- (1) *If $r(\chi) \in \{-1, 0\}$ and $\lambda = \lambda'$ and $2\lambda = p - 1$ then $\mathbf{a}_1 = 0$.*
- (2) *If $r(\chi) = 1$ and $\lambda = \lambda'$ then $\mathbf{a}_1 = 0$.*

2.9. We first look at the case where $\lambda = \lambda'$.

Proposition. *We have*

$$\text{Ext}_{U_{\chi}(W_0)}(K_{\lambda}, V_{\chi}(\lambda)) \simeq \begin{cases} K, & \text{if } \lambda \in \{0, p-1\}, \\ 0, & \text{otherwise.} \end{cases}$$

It is convenient to break up the proof of the proposition into two lemmas.

Lemma (A). *If $r(\chi) = 1$ then*

$$\text{Ext}_{U_{\chi}(W_0)}(K_{\lambda}, V_{\chi}(\lambda)) = 0.$$

Proof. Every extension of K_{λ} by $V_{\chi}(\lambda)$ can be represented by a short exact sequence of $U_{\chi}(W_0)$ -modules

$$0 \rightarrow V_{\chi}(\lambda) \rightarrow M_{\mathbf{a}} \rightarrow K_{\lambda} \rightarrow 0.$$

The claim to be proved amounts to saying that the above sequence splits, or equivalently, that $\mathbf{a} = 0$. Since all the $\mathbf{a}_j$ can be expressed as a linear combination of $\mathbf{a}_1$ and $\mathbf{a}_2$, it suffices to prove that both $\mathbf{a}_1$

and $\mathbf{a}_2$ are 0. Note that $\mathbf{a}_1 = 0$ follows from Lemma 2.8. Thus, by definition, we have

$$\mathbf{a}_j = (j-1) \left(\prod_{k=3}^j (2\lambda + k) \right) \mathbf{a}_2 \quad \text{for } 3 \leq j \leq p-2.$$

Since the height of χ is 1 it follows that $2\lambda + k \neq 0$ for all k . Therefore to prove $\mathbf{a}_2 = 0$ it suffices to show $\mathbf{a}_j = 0$ for any $2 \leq j \leq p-2$. Now, if we insert $(i, j) = (1, p-2)$ into (12) and use the fact that $(p-1)! = -1$ we get

$$-2(2\lambda - 1)\mathbf{a}_{p-2} = 0,$$

proving the claim. $\square$

Lemma (B). *If $r(\chi) \in \{-1, 0\}$ then*

$$\text{Ext}_{U_\chi(W_0)}(K_\lambda, V_\chi(\lambda)) \simeq \begin{cases} K, & \text{if } \lambda \in \{0, p-1\}, \\ 0, & \text{otherwise.} \end{cases}$$

Proof. Let $(\mathbf{a}_1, \mathbf{a}_2)$ be an arbitrary pair in K^2 and consider the vector space $M_{\mathbf{a}}$ constructed in Section 2.6. By definition, we have

$$(14) \quad \mathbf{a}_j = A_j \mathbf{a}_1 + B_j \mathbf{a}_2 \quad \text{for all } 3 \leq j \leq p-2,$$

where

$$A_j = -\frac{\lambda+1}{j-2} \left(j! + \sum_{k=4}^j (k-1)! \prod_{l=k}^j \frac{l-1}{l-3} (2\lambda + l) \right),$$

(the summation $\sum_{k=4}^j$ is understood to be 0 when $j = 3$) and

$$B_j = (j-1) \prod_{k=3}^j (2\lambda + k).$$

We rewrite (10)–(12) for the present case. First, note that (10) and (11) hold trivially since $[\lambda - \lambda - i] = p - i$ and $[\lambda - \lambda - j] = p - j$ for all $1 \leq i, j \leq p-2$. If we insert $(i, j) = (1, p-2)$ into (12) and use $(p-1)! = -1$, we see that

$$(15) \quad 2(2\lambda - 1)\mathbf{a}_{p-2} = (\lambda + 1)\mathbf{a}_1.$$

If $2 \leq i, j \leq p-2$ and $i + j \leq p$, the same formula yields

$$(16) \quad \begin{aligned} (j-i)\mathbf{a}_{i+j} &= \frac{(i+j-1)!}{(j-1)!} ((i+1)\lambda + i+j)\mathbf{a}_j \\ &\quad - \frac{(i+j-1)!}{(i-1)!} ((j+1)\lambda + i+j)\mathbf{a}_i. \end{aligned}$$

We now proceed by showing how these formulas are related. Let's begin by assuming that $\lambda \notin \{0, p-1\}$. If $2\lambda = p-1$ then Lemma 2.8 implies $\mathbf{a}_1 = 0$, which together with (14) implies

$$(17) \quad \mathbf{a}_j = (j-1)(j-1)!\mathbf{a}_2 \quad \text{for all } 3 \leq j \leq p-2.$$

It follows that $\mathbf{a}_2 = 0$ because otherwise this would mean that $\mathbf{a}_{p-2} \neq 0$ and then by (15) that $2\lambda = 1$ in contradiction with our assumption.

Suppose next that $2\lambda \neq p - 1$. If we insert $(i, j) = (2, p - 2)$ into (16), we obtain

$$(18) \quad 6\mathbf{a}_{p-2} = \mathbf{a}_2.$$

Assume first that $2\lambda = 1$. Since $\lambda \neq p - 1$ eq. (15) implies $\mathbf{a}_1 = 0$ hence (14) reduces to

$$\mathbf{a}_j = \frac{1}{6}(j-1)(j+1)\mathbf{a}_2 \quad \text{for all } 3 \leq j \leq p-2.$$

In particular, we have $2\mathbf{a}_{p-2} = \mathbf{a}_2$, which together with (18), implies $\mathbf{a}_2 = 0$, as desired.

Assume next that $2\lambda \neq 1$. Eq. (15) and (18) give the following relation

$$(19) \quad 3(\lambda + 1)\mathbf{a}_1 = (2\lambda - 1)\mathbf{a}_2,$$

which means that $\mathbf{a}_1$ and $\mathbf{a}_2$ are either both 0 or both nonzero. We assume the latter and derive a contradiction. By definition, we have $\mathbf{a}_3 = -6(\lambda + 1)\mathbf{a}_1 + 2(2\lambda + 3)\mathbf{a}_2$ hence it follows from (19) that $\mathbf{a}_3 = 8\mathbf{a}_2$. Thus, if we insert $(i, j) = (3, p - 3)$ into (16), we obtain

$$3\mathbf{a}_{p-3} = -\mathbf{a}_2.$$

Likewise, if we insert $(i, j) = (2, p - 3)$ into (16), we obtain

$$6(3\lambda - 1)\mathbf{a}_{p-3} = -(2\lambda + 1)\mathbf{a}_2.$$

Since $2\lambda + 1 \neq 0$ we can determine λ by eliminating $\mathbf{a}_2$ and $\mathbf{a}_{p-3}$

$$4\lambda = 3.$$

Note $3\mathbf{a}_{p-3} = -\mathbf{a}_2$ implies $p > 5$. If we insert $(i, j) = (2, 3)$ into (16), we see

$$\mathbf{a}_5 = (192\lambda + 360)\mathbf{a}_2 = 504\mathbf{a}_2.$$

However, by definition we have

$$\mathbf{a}_5 = \frac{1}{3}(128\lambda^2 + 656\lambda + 1080)\mathbf{a}_2 = 548\mathbf{a}_2,$$

which implies first that $p = 11$ and then that $\lambda = 9$. If we put all this together and use (14) we see $\mathbf{a}_1 = \mathbf{a}_4$. But inserting $(i, j) = (4, 6)$ into (16) implies $\mathbf{a}_4 = 0$ and thus $\mathbf{a}_1 = 0$. Contradiction!

We now move to the case where $\lambda = 0$. A necessary condition for $M_{\mathbf{a}}$ to be a $U_{\chi}(W_0)$ -module is that $2\mathbf{a}_{p-2} = -\mathbf{a}_1$, cf. (15). Since by definition

$$\mathbf{a}_{p-2} = 4\mathbf{a}_1 - \frac{3}{2}\mathbf{a}_2,$$

this implies $\mathfrak{a}_2 = 3\mathfrak{a}_1$. Now, the claim to be proved amounts to saying that each pair $(\mathfrak{a}_1, \mathfrak{a}_2)$ with $\mathfrak{a}_2 = 3\mathfrak{a}_1$ induces a $U_\chi(W_0)$ -module $M_{\mathfrak{a}}$ in the way described in Section 2.6. Using our previous notation, we have

$$A_j = -(j-2)j! \quad \text{and} \quad B_j = \frac{1}{2}(j-1)j!,$$

hence $\mathfrak{a}_2 = 3\mathfrak{a}_1$ implies

$$\mathfrak{a}_j = \frac{1}{2}(j+1)!\mathfrak{a}_1 \quad \text{for all} \quad 1 \leq j \leq p-2.$$

In particular, we have $\mathfrak{a}_{p-2} = -1/2\mathfrak{a}_1$ in consistence with (15). A straightforward computation shows that

$$\begin{aligned} (j-i)A_{i+j} &= (i+j)! \left(\frac{A_j}{(j-1)!} - \frac{A_i}{(i-1)!} \right), \\ (j-i)B_{i+j} &= (i+j)! \left(\frac{B_j}{(j-1)!} - \frac{B_i}{(i-1)!} \right). \end{aligned}$$

proving (16) for $i+j \leq p-2$. Likewise, a simple computation shows

$$\mathfrak{a}_j(p-j-2)! = \mathfrak{a}_{p-j-1}(j-1)!,$$

proving (16) for $i+j = p-1$. Since $i+j = p$ gives 0 on both sides of (16) the claim follows from Lemma 2.8.

Finally, we consider the case where $\lambda = p-1$. Here we have $A_j = 0$ and it follows from (15) that $\mathfrak{a}_{p-2} = 0$. Therefore since

$$\mathfrak{a}_j = (j-1)!\mathfrak{a}_2 \quad \text{for all} \quad 2 \leq j \leq p-2,$$

this implies first $\mathfrak{a}_2 = 0$ and then that $\mathfrak{a}_j = 0$ for all $j \geq 2$. The rest of the proof is straightforward. $\square$

2.10. Next, we address the cases where $(\lambda, \lambda') = (0, p-1)$ and $(\lambda, \lambda') = (p-1, 0)$. This can only occur if the height of χ is less than 1 because $\lambda, \lambda' \notin \mathbb{F}_p$ for $r(\chi) = 1$. The arguments presented previously would apply equally well to these cases, but we will give another, simpler, proof which relies on the fact that $\text{Ext}_{U_\chi(W_0)}(K_0, V_\chi(0))$ is 1-dimensional, cf. Proposition 2.9.

Proposition. *If $(\lambda, \lambda') = (0, p-1)$ or $(\lambda, \lambda') = (p-1, 0)$, then*

$$\text{Ext}_{U_\chi(W_0)}(K_{\lambda'}, V_\chi(\lambda)) \simeq K.$$

Proof. Let $\eta \in W^*$ be a functional of height 0. We have

$$\text{Ext}_{U_\eta(W)}(V_\eta(0), V_\eta(0)) \simeq \text{Ext}_{U_\chi(W_0)}(K_0, V_\chi(0)) \simeq K.$$

Since the reduced Verma modules $V_\eta(0)$ and $V_\eta(p-1)$ are isomorphic [2, Hilfssatz 7], we obtain

$$\text{Ext}_{U_\chi(W_0)}(K_0, V_\chi(p-1)) \simeq \text{Ext}_{U_\eta(W)}(V_\eta(0), V_\eta(p-1)) \simeq K,$$

as claimed. The other case can be handled similarly. $\square$

2.11. It is convenient to divide the case $(\lambda, \lambda') \notin \Theta(\chi)$ into two subcases depending on whether or not $\lambda + \lambda' = p - 1$. Recall that $\lambda + \lambda' = p - 1$ can only occur if $r(\chi) < 1$. The following lemma will be needed in the sequel.

Lemma. *Let i and j be two integers with $2 \leq i, j \leq p - 2$.*

- (1) *If $i > [\lambda - \lambda' - j]$ then $i + j > [\lambda - \lambda']$.*
- (2) *If $j \leq [\lambda - \lambda' - i]$ and $i + j > [\lambda - \lambda']$ then $i > [\lambda - \lambda']$.*

2.12. We denote the roots of the polynomial $x^2 - a \in K[x]$ by $\pm\sqrt{a}$.

Proposition. *If $(\lambda, \lambda') \notin \Theta(\chi)$ and $\lambda + \lambda' \neq p - 1$, then*

$$\text{Ext}_{U_\chi(W_0)}(K_{\lambda'}, V_\chi(\lambda)) \simeq \begin{cases} K, & \text{if } [\lambda - \lambda'] = 1 \text{ and } \lambda \in \{1, p - 1\}, \\ K, & \text{if } [\lambda - \lambda'] \in \{2, 3\}, \\ K, & \text{if } [\lambda - \lambda'] = 4 \text{ and } p > 5, \\ K, & \text{if } [\lambda - \lambda'] = 4 \text{ and } \lambda \in \{0, 3\}, \\ K, & \text{if } [\lambda - \lambda'] = 5 \text{ and } \lambda \in \{0, 4\}, \\ K, & \text{if } [\lambda - \lambda'] = 6 \text{ and } 2\lambda = 5 \pm \sqrt{19} \text{ and } p > 7, \\ 0, & \text{otherwise.} \end{cases}$$

Proof. Every extension of $K_{\lambda'}$ by $V_\chi(\lambda)$ can be represented by a short exact sequence of $U_\chi(W_0)$ -modules

$$(20) \quad 0 \rightarrow V_\chi(\lambda) \rightarrow M_{\mathbf{a}} \rightarrow K_{\lambda'} \rightarrow 0,$$

where $\mathbf{a}_1 = 0$ (see Lemma 2.3 B and Remark 2.3) and

$$(21) \quad \mathbf{a}_j = \frac{(-1)^j}{(j-2)!} \left(\prod_{k=2}^{j-1} (\lambda - \lambda' - k)(\lambda + \lambda' + k + 1) \right) \mathbf{a}_2, \quad 3 \leq j \leq p - 2.$$

Note that $\mathbf{a}_j = 0$ for all $j > [\lambda - \lambda']$ if $[\lambda - \lambda'] \geq 2$. This is a very useful observation which we shall use several times, often without any reference. We divide the proof into several steps depending on $[\lambda - \lambda']$.

Case 1. Suppose that $[\lambda - \lambda'] = 1$.

For the sake of simplicity, we assume that $p > 5$. Eq. (21) becomes

$$\mathbf{a}_j = \left(\prod_{k=2}^{j-1} (2\lambda + k) \right) \mathbf{a}_2 \quad \text{for all } 3 \leq j \leq p - 2.$$

If we insert $(i, j) = (2, 3)$ into (12) and then use the above formula to express $\mathbf{a}_3$ as a scalar multiple $\mathbf{a}_2$ we obtain

$$\mathbf{a}_5 = (2\lambda + 2)(18\lambda + 12)\mathbf{a}_2.$$

A necessary condition for (20) to be non-split is that $\mathfrak{a} \neq 0$, or equivalently, that $\mathfrak{a}_2 \neq 0$. If this is the case, then

$$\prod_{k=2}^4 (2\lambda + k) = (2\lambda + 2)(18\lambda + 12),$$

which implies $\lambda \in \{1, p-1\}$. (It should be noted that $\lambda = 0$ is a solution to the above equation, but it has been deliberately omitted because otherwise we would have $\lambda' = p-1$ and thus $\lambda + \lambda' = p-1$.)

Conversely, suppose that $\lambda = 1$ or $\lambda = p-1$. Since $\lambda - \lambda' = 1$, this implies $\lambda' = 0$ or $\lambda' = p-2$, respectively. Let $(\mathfrak{a}_1, \mathfrak{a}_2)$ be a pair in K^2 such that $\mathfrak{a}_1 = 0$ and consider the corresponding vector space $M_{\mathfrak{a}}$. If $\lambda = p-1$, then $\lambda + \lambda' = p-3$ hence by definition $\mathfrak{a}_j = 0$ for all $j \neq 2$ and a very simple computation shows that (10)–(12) hold. Suppose next that $\lambda = 1$. We have then

$$p-2 > [\lambda - \lambda' - 1] \quad \text{and} \quad 1 < [\lambda - \lambda' - (p-2)].$$

If we insert $(i, j) = (p-2, 1)$ into (11), we obtain zero on both sides because $(\lambda+1)+\lambda'-2 = 0$ and $\mathfrak{a}_{p-1} = 0$. Furthermore, if $2 \leq i, j \leq p-2$ then

$$[\lambda - \lambda' - i] = p+1-i \quad \text{and} \quad [\lambda - \lambda' - j] = p+1-j.$$

Thus, we may assume that

$$i \leq [\lambda - \lambda' - j] \quad \text{and} \quad j \leq [\lambda - \lambda' - i],$$

or equivalently that $i+j \leq p+1$. We have

$$\mathfrak{a}_k = \frac{1}{6}(k+1)!\mathfrak{a}_2 \quad \text{for all} \quad 2 \leq k \leq p-2,$$

which, when inserted into (12), yields

$$\begin{aligned} (j-i)\mathfrak{a}_{i+j} &= \frac{1}{6}(i+j-2)!(2i+j)(j-1)j(j+1)\mathfrak{a}_2 \\ &\quad - \frac{1}{6}(i+j-2)!(2j+i)(i-1)i(i+1)\mathfrak{a}_2. \end{aligned}$$

Since

$$(j-i)\mathfrak{a}_{i+j} = \begin{cases} \frac{1}{6}(j-i)(i+j+1)!\mathfrak{a}_2, & \text{if } i+j \leq p-2, \\ 0, & \text{if } i+j \in \{p, p \pm 1\}, \end{cases}$$

the claim follows from a straightforward computation.

Case 2. Suppose that $[\lambda - \lambda'] \in \{2, 3\}$.

Let $(\mathfrak{a}_1, \mathfrak{a}_2)$ be a pair in K^2 such that $\mathfrak{a}_1 = 0$ and consider the corresponding vector space $M_{\mathfrak{a}}$. Lemma 2.11 together with the remark following eq. (21) yields (10)–(12) for $2 \leq i, j \leq p-2$. Since furthermore $\mathfrak{a}_{p-2} = 0$ for $p > 5$ this proves (10)–(12) for all i, j in the case $p > 5$. (Note that the above reasoning works just as well for $\lambda - \lambda' = 4$.) The case $p = 5$ is straightforward and is omitted here.

Case 3. Suppose that $[\lambda - \lambda'] = 4$.

As noted above, we may assume $p = 5$ and $(i, j) \in \{(1, 3), (3, 1)\}$. We have

$$3 = [\lambda - \lambda' - 1] \quad \text{and} \quad 1 = [\lambda - \lambda' - 3].$$

Thus, if we insert $(i, j) = (1, 3)$ into (12) and then use the definition of $\mathfrak{a}_3$ to express it as a scalar multiple $\mathfrak{a}_2$, we obtain

$$2(\lambda + \lambda' + 3)(\lambda + \lambda' + 4)\mathfrak{a}_2 = 0.$$

Now, every extension of $K_{\lambda'}$ by $V_{\chi}(\lambda)$ can be represented by a short exact sequence as in (20) and such that $\mathfrak{a}_1 = 0$. The above computation shows that a necessary (and sufficient) condition for (20) to be non-split is that $\lambda + \lambda' \in \{1, 2\}$, or equivalently, that $\lambda \in \{0, 3\}$.

Case 4. Suppose that $[\lambda - \lambda'] = 5$.

First, note that the assumption implies $p > 5$ since $[\lambda - \lambda'] < p$. We consider a short exact sequence of $U_{\chi}(W_0)$ -modules as in (20). If we insert $(i, j) = (2, 3)$ into (12) and then use the definition of $\mathfrak{a}_5$ to express it as a scalar multiple $\mathfrak{a}_2$, we get

$$(22) \quad \frac{1}{6} \left(\prod_{k=3}^5 (\lambda + \lambda' + k) \right) \mathfrak{a}_2 = (2\lambda + \lambda' + 5)(\lambda + \lambda' + 3)\mathfrak{a}_2 \\ - (3\lambda + \lambda' + 5)\mathfrak{a}_2,$$

which since $\lambda + \lambda' \neq p - 1$ can be rewritten as

$$((\lambda + \lambda')(\lambda + \lambda' + 5) - 6\lambda)\mathfrak{a}_2 = 0.$$

Thus, the term inside the parentheses must equal 0 in order for (20) to be non-split. Note that the above reasoning works just as well in the case $[\lambda - \lambda'] \geq 5$. We shall make use of this later, but for now we are content to remark that in our present case this implies $\lambda \in \{0, 4\}$.

Conversely, suppose that $\lambda \in \{0, 4\}$ and let $(\mathfrak{a}_1, \mathfrak{a}_2)$ be a pair in K^2 such that $\mathfrak{a}_1 = 0$. We extend $(\mathfrak{a}_1, \mathfrak{a}_2)$ as usual to a tuple $\mathfrak{a} \in K^{p-2}$. The choice of λ together with Lemma 2.11 implies (10)–(12) for $2 \leq i, j \leq p - 2$ hence we may assume $(i, j) \in \{(1, p - 2), (p - 2, 1)\}$.

If $p = 7$ then

$$p - 2 > [\lambda - \lambda' - 1] \quad \text{and} \quad 1 > [\lambda - \lambda' - (p - 2)].$$

Since by definition $\mathfrak{a}_{p-1} = 0$, this proves (10). If $p > 7$, then

$$p - 2 > [\lambda - \lambda' - 1] \quad \text{and} \quad 1 < [\lambda - \lambda' - (p - 2)].$$

The remark following eq. (21) implies $\mathfrak{a}_{p-2} = 0$ and thus (11), thereby proving the claim.

Case 5. Suppose that $[\lambda - \lambda'] = 6$.

As previously noted, every nontrivial extension of $K_{\lambda'}$ by $V_{\chi}(\lambda)$ can be represented by a sequence of $U_{\chi}(W_0)$ -modules as in (20) and such that

$\mathfrak{a}_1 = 0$ and $(\lambda + \lambda')(\lambda + \lambda' + 5) = 6\lambda$. In our present case, this amounts to

$$\lambda = \frac{1}{2}(5 \pm \sqrt{19}).$$

For $r(\chi) < 1$ we have $\lambda \in \mathbb{F}_p$ and since $x^2 - 19$ does not split in $\mathbb{F}_7[x]$ this shows that $p > 7$. For $r(\chi) = 1$ we have $\lambda + \lambda' \notin \mathbb{F}_p$. If $p = 7$ then

$$p - 2 = [\lambda - \lambda' - 1] \quad \text{and} \quad 1 = [\lambda - \lambda' - (p - 2)].$$

But if we insert $(i, j) = (1, p - 2)$ into (12) and then express $\mathfrak{a}_{p-2}$ as a multiple of $\mathfrak{a}_2$ we obtain

$$\frac{1}{6}(\lambda + \lambda' - 1) \left(\prod_{k=2}^4 (\lambda - \lambda' - k)(\lambda + \lambda' + k + 1) \right) \mathfrak{a}_2 = 0.$$

This is only possible if $\mathfrak{a}_2 = 0$, or equivalently, if $\mathfrak{a} = 0$ contradicting the fact that the sequence (20) does not split.

Conversely, assume that $p > 7$ and let λ be as above. Let $(\mathfrak{a}_1, \mathfrak{a}_2) \in K^2$ be a pair such that $\mathfrak{a}_1 = 0$ and consider the corresponding vector space $M_{\mathfrak{a}}$. The remark following eq. (21) gives (10)–(12) for $(i, j) \in \{(1, p - 2), (p - 2, 1)\}$. By virtue of Lemma 2.11 we may assume that $i + j \leq [\lambda - \lambda']$. Since furthermore we have chosen λ in such a way that (12) holds for $(i, j) = (2, 3)$, it suffices to consider the case $(i, j) \in \{(2, 4), (4, 2)\}$. We have

$$4 = [\lambda - \lambda' - 2] \quad \text{and} \quad 2 = [\lambda - \lambda' - 4].$$

If we insert $(i, j) = (2, 4)$ into (12) and then use the definition of $\mathfrak{a}_4$ to express it as a scalar multiple $\mathfrak{a}_2$, we obtain

$$\begin{aligned} \frac{1}{12} \left(\prod_{k=3}^6 (\lambda + \lambda' + k) \right) \mathfrak{a}_2 &= \frac{1}{2} (2\lambda + \lambda' + 6) \left(\prod_{k=3}^4 (\lambda + \lambda' + k) \right) \mathfrak{a}_2 \\ &\quad - (4\lambda + \lambda' + 6) \mathfrak{a}_2, \end{aligned}$$

which gives the equation

$$(\lambda + \lambda' + 1)(\lambda + \lambda' + 6)((\lambda + \lambda')(\lambda + \lambda' + 5) - 6\lambda) \mathfrak{a}_2 = 0,$$

that holds by the definition of λ .

Case 6. Suppose that $[\lambda - \lambda'] \geq 7$.

Every extension of $K_{\lambda'}$ by $V_{\chi}(\lambda)$ can be represented by a short exact sequence of $U_{\chi}(W_0)$ -modules as in (20) and such that $\mathfrak{a}_1 = 0$. A necessary condition for (20) to be non-split is that $(\lambda + \lambda')(\lambda + \lambda' + 5) = 6\lambda$ which, in our present case, implies

$$(23) \quad \lambda' = \frac{1}{2}(-5 \pm \sqrt{24\lambda + 25}) - \lambda.$$

Assume towards contradiction that $\mathfrak{a}_2 \neq 0$. We have

$$5 \leq [\lambda - \lambda' - 2] \quad \text{and} \quad 2 \leq [\lambda - \lambda' - 5].$$

Thus, if we insert $(i, j) = (2, 5)$ into (12) and then use the definition of $\mathfrak{a}_5$ to write it as a scalar multiple $\mathfrak{a}_2$, we get

$$\frac{1}{40} \left(\prod_{k=3}^7 (\lambda + \lambda' + k) \right) \mathfrak{a}_2 = \frac{1}{6} (2\lambda + \lambda' + 7) \left(\prod_{k=3}^5 (\lambda + \lambda' + k) \right) \mathfrak{a}_2 - (5\lambda + \lambda' + 7) \mathfrak{a}_2.$$

which, together with (23), yields

$$\pm(\lambda^2 + 5\lambda + 4)\sqrt{24\lambda + 25} = 15\lambda^2 + 35\lambda + 20.$$

Squaring both sides and subtracting one from the other yields the equation

$$\frac{9}{50} \lambda(\lambda - 1)(\lambda + 1)^2(3\lambda + 2) = 0,$$

which in turn yields $\lambda \in \{0, \pm 1, -2/3\}$. Now we can insert this back into (23) to determine λ' . But first note that if

$$\lambda' = \frac{1}{2}(-5 - \sqrt{24\lambda + 25}) - \lambda,$$

then $\lambda = p - 1$ because $0, 1, -2/3$ are not solutions to

$$-(\lambda^2 + 5\lambda + 4)\sqrt{24\lambda + 25} = 15\lambda^2 + 35\lambda + 20.$$

Putting all of this together, we obtain

$$(\lambda, \lambda') \in \{(p-1, p-1), (p-1, p-2), (1, 0), (0, 0), (-2/3, -1/3)\}.$$

But none of these cases is possible since by assumption $[\lambda - \lambda'] \geq 7$ and $\lambda + \lambda' \neq p - 1$; a contradiction which can only be avoided if $\mathfrak{a}_2 = 0$. This completes the proof of the proposition. $\square$

2.13. We are left with the case where $(\lambda, \lambda') \notin \Theta(\chi)$ and $\lambda + \lambda' = p - 1$. Note that this implies $r(\chi) < 1$.

Proposition. *If $(\lambda, \lambda') \notin \Theta(\chi)$ and $\lambda + \lambda' = p - 1$, then*

$$\text{Ext}_{U_\chi(W_0)}(K_{\lambda'}, V_\chi(\lambda)) \simeq \begin{cases} K, & \text{if } [\lambda - \lambda'] \in \{2, 3, 4\}, \\ K, & \text{if } [\lambda - \lambda'] = 6 \text{ and } p = 19, \\ 0, & \text{otherwise.} \end{cases}$$

Proof. Let $(\lambda, \lambda') \notin \Theta(\chi)$ such that $\lambda + \lambda' = p - 1$. Every extension of $K_{\lambda'}$ by $V_\chi(\lambda)$ can be represented by a short exact sequence of $U_\chi(W_0)$ -modules as (20). The assumption $\lambda + \lambda' = p - 1$ implies $\mathfrak{a}_2 = 0$, see Lemma 2.3 B and Remark 2.3. Using the foregoing notation, we have

$$A_j = \frac{(-1)^j}{j-2} \left(\prod_{k=0}^{j-2} (2\lambda - k) \right) \left((j-2)\lambda - 1 + j + \sum_{k=4}^j \frac{(j-k)!}{(j-3)!} \cdot ((j+1-k)\lambda + (j+2-k)) \prod_{l=1}^{k-3} (j-l) \right).$$

(The summation $\sum_{k=4}^j$ is understood to be 0 when $j = 3$.) Note that $\mathfrak{a}_j = 0$ for all $j > [\lambda - \lambda']$. Furthermore, it should also be noted that $\lambda - \lambda' \neq 1$ since otherwise we would have $(\lambda, \lambda') = (0, p-1)$ in contradiction to the assumption that $(\lambda, \lambda') \notin \Theta(\chi)$.

Case 1. Suppose that $[\lambda - \lambda'] \in \{2, 3, 4\}$.

Conversely, every pair $(\mathfrak{a}_1, \mathfrak{a}_2) \in K^2$ with $\mathfrak{a}_2 = 0$ gives rise to a module $M_{\mathfrak{a}}$ as described in Section 2.6: Lemma 2.11 yields (10)–(12) for $2 \leq i, j \leq p-2$. For $(i, j) \in \{(1, p-2), (p-2, 1)\}$ the computations are simple and are omitted for brevity.

Case 2. Suppose that $[\lambda - \lambda'] = 5$.

Consider a short exact sequence of $U_{\chi}(W_0)$ -modules as in (20). If we insert $(i, j) = (2, 3)$ into (12) and then insert the expressions $\mathfrak{a}_3 = A_3\mathfrak{a}_1$ and $\mathfrak{a}_5 = A_5\mathfrak{a}_1$ into the result we get

$$\lambda(\lambda+2)(\lambda+4) \left(\prod_{k=1}^3 (2\lambda - k) \right) \mathfrak{a}_1 = \frac{1}{3} \lambda (13\lambda + 22) \left(\prod_{k=1}^3 (2\lambda - k) \right) \mathfrak{a}_1.$$

Now, the assumptions $\lambda + \lambda' = p-1$ and $(\lambda, \lambda') \notin \Theta(\chi)$ imply $\lambda \neq 0$. Since, in addition, $[\lambda - \lambda'] = 5$, the above equation reduces to

$$3(\lambda+2)(\lambda+4)\mathfrak{a}_1 = (13\lambda+22)\mathfrak{a}_1,$$

which, by subtracting the right side from the left side, gives

$$(3\lambda^2 + 5\lambda + 2)\mathfrak{a}_1 = 0.$$

We get a quadratic equation which can be solved easily; the roots are $p-1$ and $-2/3$. But, we can immediately exclude $\lambda = p-1$ since this would imply $(\lambda, \lambda') = (p-1, 0)$. All the above reasoning works just as well in the case where $[\lambda - \lambda'] \geq 5$. We shall make use of this later, but for now we are content to remark that in our present case $\lambda + \lambda' = p-1$ implies $(\lambda, \lambda') = (-2/3, -1/3)$ contradicting the fact that $\lambda - \lambda' = 5$.

Case 3. Suppose that $[\lambda - \lambda'] = 6$.

As noted previously, a necessary condition for (20) to be non-split is that $\lambda = -2/3$. The assumption $\lambda + \lambda' = p-1$ implies $\lambda - \lambda' = 2\lambda + 1$ which since $\lambda - \lambda' = 6$ means $\lambda = 5/2$ and hence $p = 19$. Conversely, let $p = 19$ and $(\lambda, \lambda') = (12, 6)$. (Note $12 = 5/2$ in characteristic $p = 19$.) Let $(\mathfrak{a}_1, \mathfrak{a}_2)$ be a pair in K^2 such that $\mathfrak{a}_2 = 0$ and consider the corresponding vector space $M_{\mathfrak{a}}$. Eq. (10)–(12) follow immediately for $(i, j) \in \{(1, p-2), (p-2, 1)\}$ since $\mathfrak{a}_{p-2} = 0$. By virtue of Lemma 2.11 we may assume that $(i, j) \in \{(2, 4), (4, 2)\}$. We have

$$2 = [\lambda - \lambda' - 4] \quad \text{and} \quad 4 = [\lambda - \lambda' - 2].$$

If we insert $(i, j) = (2, 4)$ into (12) and then insert the expressions $\mathfrak{a}_4 = A_4 \mathfrak{a}_1$ and $\mathfrak{a}_6 = A_6 \mathfrak{a}_1$ into the result, we get

$$\frac{1}{3} \lambda (77\lambda + 125) \left(\prod_{k=1}^4 (2\lambda - k) \right) \mathfrak{a}_1 = \lambda (\lambda + 5) (5\lambda + 9) \left(\prod_{k=1}^4 (2\lambda - k) \right) \mathfrak{a}_1.$$

Keeping in mind that the characteristic is 19, a very simple computation shows that the above equation holds for all $\mathfrak{a}_1$.

Case 4. Suppose that $[\lambda - \lambda'] \geq 7$.

Suppose we are given a short exact sequence of $U_\chi(W_0)$ -modules as in (20) and such that $\mathfrak{a}_2 = 0$ and $\lambda = -2/3$. We have

$$2 \leq [\lambda - \lambda' - 5] \quad \text{and} \quad 5 \leq [\lambda - \lambda' - 2].$$

If we insert $(i, j) = (2, 5)$ into (12), we obtain

$$\begin{aligned} \frac{6}{10} \lambda (87\lambda + 137) \left(\prod_{k=1}^5 (2\lambda - k) \right) \mathfrak{a}_1 &= \frac{2}{3} \lambda (\lambda + 6) (13\lambda + 22) \\ &\cdot \left(\prod_{k=1}^5 (2\lambda - k) \right) \mathfrak{a}_1. \end{aligned}$$

Since $\lambda \neq 0$ and $[\lambda - \lambda'] \geq 7$, the above equation reduces to

$$9(87\lambda + 137) \mathfrak{a}_1 = 10(\lambda + 6)(13\lambda + 22) \mathfrak{a}_1,$$

which, by subtracting the left side from the right side, gives

$$(130\lambda^2 + 217\lambda + 87) \mathfrak{a}_1 = 0.$$

The claim follows since $-2/3$ is not a root of the polynomial inside the brackets. $\square$

2.14. We summarize the preceding results as a theorem as follows.

Theorem. *We have the following three cases*

(1) *If $\lambda' \in \{0, p-1\}$, then*

$$\text{Ext}_{U_\chi(W)}(V_\chi(\lambda'), V_\chi(\lambda)) \simeq \begin{cases} K, & \text{if } \lambda \in \{0, 1, 2, 3, 4, p-1\}, \\ 0, & \text{otherwise.} \end{cases}$$

(2) *If $\lambda \in \{0, p-1\}$, then*

$$\text{Ext}_{U_\chi(W)}(V_\chi(\lambda'), V_\chi(\lambda)) \simeq \begin{cases} K, & \text{if } \lambda' \in \{0\} \cup \{p-i \mid 1 \leq i \leq 5\}, \\ 0, & \text{otherwise.} \end{cases}$$

(3) If $\lambda, \lambda' \notin \{0, p-1\}$, then

$$\begin{aligned} & \text{Ext}_{U_\chi(W)}(V_\chi(\lambda'), V_\chi(\lambda)) \\ & \simeq \begin{cases} K, & \text{if } [\lambda - \lambda'] \in \{2, 3\}, \\ K, & \text{if } [\lambda - \lambda'] = 4 \text{ and } p \neq 5, \\ K, & \text{if } [\lambda - \lambda'] = 6 \text{ and } 2\lambda = 5 \pm \sqrt{19} \text{ and } p > 7, \\ 0, & \text{otherwise.} \end{cases} \end{aligned}$$

3. HEIGHT -1

Throughout this section we will assume that $r(\chi) = -1$, or equivalently, that $\chi = 0$. Furthermore, we let as usual λ and λ' be elements in $\Lambda(0) \simeq \mathbb{F}_p$.

3.1. $V_0(\lambda)$ and $V_0(\lambda')$. This case is fully described in Section 2. The reader is referred to Theorem 2.14.

3.2. $V_0(\lambda)$ and K . Our approach will follow the one taken earlier in Section 2. We have an isomorphism of vector spaces

$$\text{Ext}_{U_0(W)}(V_0(\lambda), K) \simeq \text{Ext}_{U_0(W_0)}(K_\lambda, K),$$

where K denotes the trivial W -module. Suppose that we have a short exact sequence of $U_0(W_0)$ -modules

$$(24) \quad 0 \rightarrow K \xrightarrow{f} M \xrightarrow{g} K_\lambda \rightarrow 0,$$

and fix bases v and v' for K and K_λ , respectively. Let $\{w, w'\}$ be a basis for M such that $f(v) = w$ and $g(w') = v'$. Since $U_0(Ke_0)$ is semisimple we may choose w' such that $e_0 w' = \lambda w'$. Note that the λ weight space in M is one dimensional so w' is unique with these properties. Now, we have clearly $e_i w = 0$ for all i . Furthermore, if $e_1 w'$ and $e_2 w'$ are both 0, then $e_i w' = 0$ for all $i > 0$. This can be seen by a simple induction argument since

$$(i-1)e_{i+1}w' = (e_1e_i - e_ie_1)w'.$$

The above observations give a necessary condition for (24) to be non-split, namely, that $e_1 w' \neq 0$ or $e_2 w' \neq 0$. We can interpret this condition in terms of λ . Indeed, since $e_i w'$ belongs to the weight space $M_{\lambda+i}$, we have $e_i w' = 0$ if $\lambda + i \neq 0$ and $i > 0$. Thus $e_1 w' = 0$ and $e_2 w' = 0$ for $\lambda \notin \{p-1, p-2\}$. This leads to the following lemma

Lemma. *If $\lambda \notin \{p-1, p-2\}$ then*

$$\text{Ext}_{U_0(W_0)}(K_\lambda, K) = 0.$$

We consider the cases $\lambda \in \{p-1, p-2\}$ separately. Note that $\lambda = p-1$ is not interesting for our purpose because $V_0(p-1)$ is not simple; we shall, nevertheless, include it in our study for the sake of

completeness. Fix $\lambda = p - k$ for $k \in \{1, 2\}$. There exists $\mathfrak{a} \in K$ such that for every $i > 0$

$$e_i w' = \begin{cases} \mathfrak{a}w, & \text{if } i = k, \\ 0, & \text{otherwise.} \end{cases}$$

As in Section 2, we construct a well-defined injective homomorphism

$$\text{Ext}_{U_0(W_0)}(K_\lambda, K) \rightarrow K,$$

which sends the class of (24) to $\mathfrak{a}$. Conversely, for each $\mathfrak{a} \in K$ consider a short exact sequence of vector spaces

$$0 \rightarrow K \xrightarrow{f} M_{\mathfrak{a}} \xrightarrow{g} K_\lambda \rightarrow 0,$$

and choose a basis $\{w, w'\}$ for $M_{\mathfrak{a}}$ such that $f(v) = w$ and $g(w') = v'$ where, as before, v and v' are bases for K and K_λ , respectively. For each $0 \leq i \leq p - 2$ define an endomorphism $E_i \in \text{End}_K(M_{\mathfrak{a}})$ of $M_{\mathfrak{a}}$ such that $E_i w = 0$ for all i and

$$E_i w' = \begin{cases} \lambda w', & \text{if } i = 0, \\ \mathfrak{a}w, & \text{if } i = k, \\ 0, & \text{otherwise.} \end{cases}$$

Furthermore, set $E_i^{[p]} = \delta_{i0} E_i$ for all $i = 0, 1, \dots, p - 2$ and $E_i = 0$ for $i \neq 0, 1, \dots, p - 2$. We have

$$(E_i^p - E_i^{[p]})w' = 0 \quad \text{for all } 0 \leq i \leq p - 2,$$

and

$$[E_i, E_j]w' = (j - i)E_{i+j}w' \quad \text{for all } i \text{ and } j.$$

To prove the above formula one can assume $i \in \{0, k\}$ and that $i \neq j$ since otherwise one would obtain 0 on both sides of the equation. We summarize the results in this section into the following Proposition

Proposition. *We have*

$$\text{Ext}_{U_0(W)}(V_0(\lambda), K) \simeq \begin{cases} K, & \text{if } \lambda \in \{p - 1, p - 2\}, \\ 0, & \text{otherwise.} \end{cases}$$

Evidently, K is a self-dual module in the sense that it is isomorphic to its dual. Thanks to the isomorphism

$$\text{Ext}_{U_0(W)}(K, V_0(\lambda)) \simeq \text{Ext}_{U_0(W)}(V_0(\lambda)^*, K^*),$$

and the fact that $V_0(\lambda)^*$ and $V_0(p - 1 - \lambda)$ are isomorphic we obtain

Corollary. *We have*

$$\text{Ext}_{U_0(W)}(K, V_0(\lambda)) \simeq \begin{cases} K, & \text{if } \lambda = \{0, 1\}, \\ 0, & \text{otherwise.} \end{cases}$$

3.3. $V_0(\lambda)$ and S . We proceed to describe $\text{Ext}_{U_0(W)}(V_0(\lambda), S)$ where S denotes the $(p-1)$ -dimensional simple $U_0(W)$ -module. We have an isomorphism

$$\text{Ext}_{U_0(W)}(V_0(\lambda), S) \simeq \text{Ext}_{U_0(W_0)}(K_\lambda, S).$$

Thus, classifying the extensions of $V_0(\lambda)$ by S reduces to classifying the extensions of K_λ by S . Suppose that we have a short exact sequence of $U_0(W_0)$ -modules

$$(25) \quad 0 \rightarrow S \xrightarrow{f} M \xrightarrow{g} K_\lambda \rightarrow 0,$$

and fix bases $v_0, v_1, \dots, v_{p-2}$ and v' of S and K_λ , respectively. Let

$$(26) \quad w_0, \dots, w_{p-2}, w'$$

be a basis of M such that $f(v_i) = w_i$ for all i and $g(w') = v'$; we may choose w' such that $e_0 w' = \lambda w'$. We have

$$e_j w_i = \begin{cases} (-1)^{j+1} \frac{(i+1)!}{(i-j)!} w_{i-j}, & \text{if } j \leq i, \\ 0, & \text{otherwise.} \end{cases}$$

By weight considerations there exists $\mathfrak{a}_j \in K$ ($j > 0$) such that

$$e_j w' = \begin{cases} \mathfrak{a}_j w_{[-\lambda-j-1]}, & \text{if } \lambda + j \neq 0, \\ 0, & \text{otherwise.} \end{cases}$$

Furthermore, we set

$$(27) \quad \mathfrak{a}_j = 0 \quad \text{if } \lambda + j = 0 \quad \text{or } j \notin \{1, 2, \dots, p-2\},$$

For $\lambda \neq 0$ the λ weight space in M is two-dimensional; it is generated by w' and $w_{[-\lambda-1]}$ so any different choice for w' has the form $w' + bw_{[-\lambda-1]}$ for some $b \in K$. Obviously, this leads to the same $e_j w'$ if $e_j w_{[-\lambda-1]} = 0$. In particular, we see that all the $\mathfrak{a}_j$ are determined by M if $\lambda = p-1$. The same holds for $\lambda = 0$ since w' is unique in this case; we have $M_0 = Kw'$.

Lemma (A). *If $\lambda \in \{0, p-1\}$ then all the $\mathfrak{a}_j$ are determined by M .*

For $\lambda \notin \{0, p-1\}$ we have $e_1 w_{[-\lambda-1]} \neq 0$. Thus, the discussion preceding Lemma A shows that we can choose w' uniquely such that $e_1 w' = 0$.

Lemma (B). *If $\lambda \notin \{0, p-1\}$ then there is a unique choice for w' such that $e_1 w' = 0$.*

We will henceforth always assume that w' is the unique choice from Lemma B if $\lambda \notin \{0, p-1\}$. As in Section 2 we obtain an injective homomorphism $\text{Ext}_{U_0(W_0)}(K_\lambda, S) \rightarrow K^2$ which sends the class of (25) to $(\mathfrak{a}_1, \mathfrak{a}_2) \in K^2$ and such that $\mathfrak{a}_1 = 0$ for $\lambda \notin \{0, p-1\}$. We have

$$(28) \quad \mathfrak{a}_j = A_j \mathfrak{a}_1 + B_j \mathfrak{a}_2 \quad \text{for all } 3 \leq j \leq p-2,$$

where

$$A_j = -\frac{1}{j-2} \left(\prod_{k=1}^j (\lambda + k) \right) \left(1 + \sum_{k=4}^j \frac{(j-k)!}{(j-3)!} \prod_{l=1}^{k-3} (\lambda + j - l) \right),$$

(the summation $\sum_{k=4}^j$ is understood to be 0 when $j = 3$) and

$$B_j = \frac{(-1)^j}{(j-2)!} \prod_{k=2}^{j-1} (-\lambda - 1 - k)(\lambda + k).$$

Conversely, let $(\mathbf{a}_1, \mathbf{a}_2)$ be a pair in K^2 such that, cf. (27)

$$\mathbf{a}_1 = 0 \quad \text{if} \quad \lambda = p-1,$$

$$\mathbf{a}_2 = 0 \quad \text{if} \quad \lambda = p-2.$$

We extend $(\mathbf{a}_1, \mathbf{a}_2)$ to a tuple $\mathbf{a} = (\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_{p-2}) \in K^{p-2}$ by using (28) and consider a short exact sequence

$$(29) \quad 0 \rightarrow S \xrightarrow{f} M_{\mathbf{a}} \xrightarrow{g} K_{\lambda} \rightarrow 0,$$

where $M_{\mathbf{a}}$ is a vector space with a basis as in (26). For each $0 \leq j \leq p-2$ we define an endomorphism $E_j \in \text{End}_K(M_{\mathbf{a}})$ such that

$$E_j w_i = \begin{cases} (-1)^{j+1} \frac{(i+1)!}{(i-j)!} w_{i-j}, & \text{if } j \leq i, \\ 0, & \text{otherwise,} \end{cases}$$

and $E_0 w' = \lambda w'$ and for every $1 \leq j \leq p-2$

$$E_j w' = \begin{cases} \mathbf{a}_j w_{[-\lambda-j-1]}, & \text{if } \lambda + j \neq 0, \\ 0, & \text{otherwise.} \end{cases}$$

Furthermore, we set $E_j^{[p]} = \delta_{j0} E_j$ for all j and $E_j = 0$ for $j \notin \{0, 1, \dots, p-2\}$. We have

$$(E_j^p - E_j^{[p]})w' = 0 \quad \text{for all } j.$$

Except for the change in notation, the formula $[E_i, E_j]w' = (j-i)E_{i+j}w'$ leads to the same equations as in Section 2.7; there is no need to include them here.

Proposition. *We have*

$$\text{Ext}_{U_0(W)}(V_0(\lambda), S) \simeq \begin{cases} K, & \text{if } \lambda \in \{0\} \cup \{p-i \mid 3 \leq i \leq 5\}, \\ 0, & \text{otherwise.} \end{cases}$$

Proof. The short exact sequence

$$(30) \quad 0 \rightarrow S \rightarrow V_0(0) \xrightarrow{\pi} K \rightarrow 0$$

induces the long exact sequence of vector spaces

$$(31) \quad \begin{aligned} 0 \rightarrow \text{Hom}_{U_0(W)}(V_0(\lambda), S) &\rightarrow \text{Hom}_{U_0(W)}(V_0(\lambda), V_0(0)) \\ &\rightarrow \text{Hom}_{U_0(W)}(V_0(\lambda), K) \rightarrow \text{Ext}_{U_0(W)}(V_0(\lambda), S) \\ &\rightarrow \text{Ext}_{U_0(W)}(V_0(\lambda), V_0(0)) \rightarrow \text{Ext}_{U_0(W)}(V_0(\lambda), K) \rightarrow \dots \end{aligned}$$

Due to Schur's Lemma, we have $\text{Hom}_{U_0(W)}(V_0(\lambda), K) = 0$ for $\lambda \notin \{0, p-1\}$. We obtain, in these cases, the exact sequence

$$(32) \quad 0 \rightarrow \text{Ext}_{U_0(W)}(V_0(\lambda), S) \rightarrow \text{Ext}_{U_0(W)}(V_0(\lambda), V_0(0)) \\ \rightarrow \text{Ext}_{U_0(W)}(V_0(\lambda), K) \rightarrow \cdots$$

Now $\text{Hom}_{U_0(W)}(V_0(0), K)$ is 1-dimensional and generated by the surjection π in (30). The map $\text{Hom}_{U_0(W)}(V_0(0), V_0(0)) \rightarrow \text{Hom}_{U_0(W)}(V_0(0), K)$ appearing in (31) is surjective because it maps the identity $\text{id}_{U_0(W)}$ on $U_0(W)$ to π . Thus, we obtain an exact sequence as in (32)

$$0 \rightarrow \text{Ext}_{U_0(W)}(V_0(0), S) \rightarrow \text{Ext}_{U_0(W)}(V_0(0), V_0(0)) \\ \rightarrow \text{Ext}_{U_0(W)}(V_0(0), K) \rightarrow \cdots$$

Consequently, we have for every $\lambda \neq p-1, p-2$

$$\text{Ext}_{U_0(W)}(V_0(\lambda), S) \simeq \text{Ext}_{U_0(W)}(V_0(\lambda), V_0(0)),$$

which, by Proposition 2.14, implies

$$\text{Ext}_{U_0(W)}(V_0(\lambda), S) \simeq \begin{cases} K, & \text{if } \lambda \in \{0\} \cup \{p-i \mid 3 \leq i \leq 5\}, \\ 0, & \text{if } \lambda \notin \{0\} \cup \{p-i \mid 1 \leq i \leq 5\}. \end{cases}$$

Next, suppose that $\lambda \in \{p-1, p-2\}$. We prove that every short exact sequence of $U_0(W)$ -modules as in (29) splits, or equivalently, that $\mathbf{a}_1 = 0$ and $\mathbf{a}_2 = 0$. The case $\lambda = p-2$ comes almost at once from the observations preceding the proposition; indeed $\mathbf{a}_1 = 0$ because of Lemma B and $\mathbf{a}_2 = 0$ by definition, see (27). For $\lambda = p-1$ it follows from (27) that $\mathbf{a}_1 = 0$. To prove $\mathbf{a}_2 = 0$ note first that (28) implies $\mathbf{a}_2 = -2\mathbf{a}_{p-2}$. However, since

$$1 \leq [-1 - \lambda - (p-2)] \quad \text{and} \quad p-2 \leq [-1 - \lambda - 1],$$

we see by inserting $(i, j) = (1, p-2)$ into (12) that $\mathbf{a}_{p-2} = 0$. This is only possible if $\mathbf{a}_2 = 0$ so the proposition is proved. $\square$

The module S is a self-dual module. Thanks to the isomorphism

$$\text{Ext}_{U_0(W)}(S, V_0(\lambda)) \simeq \text{Ext}_{U_0(W)}(V_0(\lambda)^*, S^*)$$

and the fact that $V_0(\lambda)^*$ and $V_0(p-1-\lambda)$ are isomorphic we obtain

Corollary. *We have*

$$\text{Ext}_{U_0(W)}(S, V_0(\lambda)) \simeq \begin{cases} K, & \text{if } \lambda \in \{2, 3, 4, p-1\}, \\ 0, & \text{otherwise.} \end{cases}$$

3.4. Self-extensions of K . There are no nontrivial self-extensions of the trivial module K .

Proposition. *We have*

$$\mathrm{Ext}_{U_0(W)}(K, K) = 0$$

Proof. Suppose we have a short exact sequence of $U_0(W)$ -modules

$$(33) \quad 0 \rightarrow K \xrightarrow{f} M \xrightarrow{g} K \rightarrow 0,$$

and let $\{w, w'\}$ be a basis of M such that $Kw = \mathrm{img} f$. We have clearly $Ww = 0$. Furthermore, $Ww' \subset Kw$ since $Ww' \subset \ker g$. Thus, every $x \in W$ acting on M can be represented by a matrix

$$\begin{bmatrix} 0 & \phi(x) \\ 0 & 0 \end{bmatrix},$$

where $\phi : W \rightarrow K$ is a homomorphism of Lie algebras. That ϕ preserves the Lie algebra structure follows from the fact that M is a W -module and hence that $W \rightarrow \mathfrak{gl}(M) \simeq \mathfrak{gl}(2, K)$ is a homomorphism of Lie algebras. Now, if (33) is non-split, then $\phi(x) \neq 0$ for some $x \in W$. The kernel of ϕ is then an ideal of codimension 1, in apparent contradiction with the fact that W is simple. $\square$

3.5. Self-extensions of S . There are no nontrivial self-extensions of the module S .

Proposition. *We have*

$$\mathrm{Ext}_{U_0(W)}(S, S) = 0.$$

Proof. The short exact sequence

$$0 \rightarrow S \rightarrow V_0(0) \rightarrow K \rightarrow 0$$

induces the long exact sequence

$$\begin{aligned} 0 &\rightarrow \mathrm{Hom}_{U_0(W)}(S, S) \rightarrow \mathrm{Hom}_{U_0(W)}(S, V_0(0)) \\ &\rightarrow \mathrm{Hom}_{U_0(W)}(S, K) \rightarrow \mathrm{Ext}_{U_0(W)}(S, S) \\ &\rightarrow \mathrm{Ext}_{U_0(W)}(S, V_0(0)) \rightarrow \cdots, \end{aligned}$$

which by Corollary 3.3 and Schur's lemma implies the claim. $\square$

3.6. S and K . The proof of the next proposition makes use of the fact that $\mathrm{Hom}_{U_0(W)}(V_0(0), S) = 0$.

Proposition. *We have*

$$\mathrm{Ext}_{U_0(W)}(K, S) \simeq K^2.$$

Proof. The short exact sequence

$$0 \rightarrow S \rightarrow V_0(0) \rightarrow K \rightarrow 0$$

induces the long exact sequence

$$\begin{aligned} 0 \rightarrow \operatorname{Hom}_{U_0(W)}(K, S) &\rightarrow \operatorname{Hom}_{U_0(W)}(V_0(0), S) \\ &\rightarrow \operatorname{Hom}_{U_0(W)}(S, S) \rightarrow \operatorname{Ext}_{U_0(W)}(K, S) \\ &\rightarrow \operatorname{Ext}_{U_0(W)}(V_0(0), S) \rightarrow \operatorname{Ext}_{U_0(W)}(S, S) \rightarrow \cdots, \end{aligned}$$

which in turn induces the exact sequence

$$0 \rightarrow \operatorname{Hom}_{U_0(W)}(S, S) \rightarrow \operatorname{Ext}_{U_0(W)}(K, S) \rightarrow \operatorname{Ext}_{U_0(W)}(V_0(0), S) \rightarrow 0.$$

The claim follows from Schur's lemma and Proposition 3.3. $\square$

Since K and S are self-dual modules, the isomorphism

$$\operatorname{Ext}_{U_0(W)}(K, S) \simeq \operatorname{Ext}_{U_0(W)}(S^*, K^*)$$

implies the following corollary

Corollary. *We have*

$$\operatorname{Ext}_{U_0(W)}(S, K) \simeq K^2.$$

4. HEIGHT 0 AND 1

All the work has been done in Section 2; Theorem 2.14 gives a complete classification of the χ -reduced Verma modules having character χ of height 0 and 1.

5. HEIGHT $p - 1$

Recall that every simple $U_\chi(W)$ -module with one exception is projective. The remaining simple module L has a projective cover with two composition factors both isomorphic to L . We have a short exact sequence

$$0 \rightarrow L \rightarrow P \rightarrow L \rightarrow 0,$$

where P is a projective module. This induces the long exact sequence

$$\begin{aligned} 0 \rightarrow \operatorname{Hom}_{U_\chi(W)}(L, L) &\rightarrow \operatorname{Hom}_{U_\chi(W)}(P, L) \\ &\rightarrow \operatorname{Hom}_{U_\chi(W)}(L, L) \rightarrow \operatorname{Ext}_{U_\chi(W)}(L, L) \\ &\rightarrow \operatorname{Ext}_{U_\chi(W)}(P, L) \rightarrow \cdots, \end{aligned}$$

which in turn implies $\operatorname{Ext}_{U_\chi(W)}(L, L) \simeq K$. We have therefore

Proposition. *If M and N are two simple $U_\chi(W)$ -modules then*

$$\operatorname{Ext}_{U_\chi(W)}(M, N) \simeq \begin{cases} K, & \text{if } N = M = L, \\ 0, & \text{otherwise.} \end{cases}$$

REFERENCES

- [1] B. D. Boe, D. K. Nakano, and E. Wiesner. Ext^1 -quivers for the Witt algebra $W(1, 1)$. *Journal of Algebra*, 322:1548–1564, 2009.
- [2] H. J. Chang. Über Wittsche Lie-Ringe. *Abh. Math. Sem. Univ. Hamburg*, 14:151–184, 1941.
- [3] J. Feldvoss and D. K. Nakano. Representation Theory of the Witt Algebra. *J. Algebra*, 203:447–469, 1998.

DEPARTMENT OF MATHEMATICS, AARHUS UNIVERSITY, BUILDING 530, NY
MUNKEGADE, 8000 AARHUS C, DENMARK

E-mail address: khalid@imf.au.dk